

## Genetic testing presents the opportunity for physicians to potentially identify a causal mutation in their patients.

In the past, genetic testing was able to pinpoint a mutation as contributing to epileptic encephalopathies in only a handful of cases.<sup>1,2</sup> However, today this frequency has risen to an estimated 15-20%<sup>3</sup> and is expected to increase further as new genes are discovered in the field.

## One question remaining in the field is this: **When is it most valuable to order genetic testing for people with epilepsy?**

The International League Against Epilepsy (ILAE) — an association of healthcare professionals dedicated to improving epilepsy education and research — has provided some perspective on this topic. This group advises that physicians consider prescribing genetic testing when there is:

- **Clinical validity** (the physician has made an informed opinion that their patient has epilepsy or a seizure disorder with a genetic basis), and
- **Clinical utility** (the results of the test are likely to positively impact the patient’s health or well-being).<sup>4</sup>

It’s important to know that if clinical utility and clinical validity can be established for genetic testing, insurers may cover the cost of testing.

While the ultimate decision of whether to order genetic testing — particularly in regard to clinical validity — rests with the treating physician, there are a number of instances where the clinical utility of genetic testing has been demonstrated. Examples are shown in the table below.

This identification of *specific genetic mutations* may have implications for clinical management by enabling an earlier diagnosis, informing therapy choice, or preventing unnecessary invasive procedures.

| Gene                  | Clinical utility of testing for mutation  |
|-----------------------|---|
| <i>ALDH7A1</i>        | Dietary lysine restriction along with pyridoxine treatment may reduce seizures and allow for developmental gains. <sup>5</sup>                                    |
| <i>DEPDC5</i>         | Surgery has been shown to reduce seizures in some patients with mutations in this gene. <sup>6</sup>  |
| <i>EMP2A or EMP2B</i> | Sodium channel blockers and GABAergic drugs may exacerbate seizures. <sup>7,8</sup>   |
| <i>FOLR</i>           | Patients with mutations in <i>FOLR</i> may respond well to treatment with folic acid. <sup>9</sup> Treatment with folic acid may worsen symptoms. <sup>7,10</sup> |
| <i>GRIN2A</i>         | Memantine has been shown to reduce seizures in those with specific <i>GRIN2A</i> mutations. <sup>11</sup>   |
| <i>KCNT1</i>          | Treatment with quinidine may reduce seizures in patients with specific <i>KCNT1</i> mutations. <sup>12,13,14</sup>  |
| <i>PNPO</i>           | Pyridoxal-5-phosphate treatment can reduce seizure recurrence and may improve cognitive function. <sup>15</sup>   |
| <i>POLG</i>           | Treatment with valproic acid may induce liver failure in patients with <i>POLG</i> deficiency. <sup>16</sup>  |
| <i>PRRT2</i>          | Seizures induced by <i>PRRT2</i> mutations may respond well to carbamazepine, phenobarbital, valproate, or zonisamide. <sup>7,17,18</sup>                         |
| <i>SCN1A</i>          | Carbamazepine and phenytoin may exacerbate seizures. <sup>2,7,19</sup>  |
| <i>SCN8A</i>          | Carbamazepine and phenytoin may help reduce seizures in patients with specific <i>SCN8A</i> mutations. <sup>20,21,22</sup>  |
| <i>SLC2A1</i>         | A ketogenic diet can reduce seizures in these patients. <sup>2</sup> Steroids and carbonic anhydrase inhibitors may improve seizure control. <sup>7</sup>         |
| <i>TSC1 or TSC2</i>   | Monitoring of certain organs in these patients may be essential. <sup>23</sup> Treatment with everolimus may reduce seizures. <sup>24</sup>                       |

**Pairnomix empowers people living with rare disease by performing personalized genetic research that enables physicians to make better-informed healthcare decisions. We're here to answer your questions about genetic research.**

**Contact us to learn more.**

*Pairnomix helps people with epilepsy who have already received genetic sequencing and have successfully identified the gene mutation that is known or believed to cause their condition. We do not perform genetic sequencing tests, but we encourage you to speak with your physician about your options.*

- <sup>1</sup>Thomas RH, Berkovic SF. The hidden genetics of epilepsy-a clinically important new paradigm. *Nat Rev Neurol.* 2014;10:283.
- <sup>2</sup>Poduri A, Sheidley BR, Shostak S, et al. Genetic testing in the epilepsies-developments and dilemmas. *Nat Rev Neurol.* 2014;10:293.
- <sup>3</sup>EpiPM Consortium. A roadmap for precision medicine in the epilepsies. *Lancet Neurol.* 2015;14:1219.
- <sup>4</sup>Ottman R, Hirose S, Jain S, et al. Genetic testing in the epilepsies- report of the ILAE Genetics Commission. *Epilepsia.* 2010;51(4):655.
- <sup>5</sup>Van Karnebeek CD, Hartmann H, Jaggumantri S, et al. Lysine restricted diet for pyridoxine-dependent epilepsy: first evidence and future trials. *Mol Genet Metab.* 2012;107:335.
- <sup>6</sup>Baulac S, Ishida S, Marsan E, et al. Familial focal epilepsy with focal cortical dysplasia due to *DEPDC5* mutations. *Ann Neurol.* 2015;77:675.
- <sup>7</sup>Ream MA, Patel AD. Obtaining genetic testing in pediatric epilepsy. *Epilepsia.* 2015;56(10):1505.
- <sup>8</sup>Medina MT, Martínez-Juárez IE, Durón RM et al. Treatment of myoclonic epilepsies of childhood, adolescence, and adulthood. *Adv Neurol.* 2005;95:307.
- <sup>9</sup>Surtees R, Wolf N. Treatable neonatal epilepsy. *Arch Dis Child.* 2007;92:659.
- <sup>10</sup>Wu D, Partridge WM. Blood-brain barrier transport of reduced folic acid. *Pharm Res.* 1999;16:415.
- <sup>11</sup>Pierson TM, Yuan H, Marsh ED, et al. *GRIN2A* mutation and early-onset epileptic encephalopathy: personalized therapy with memantine. *Ann Clin Trans Neurol.* 2014;1(3):190.
- <sup>12</sup>Bearden D, Strong A, Ehn J, et al. Targeted treatment of migrating partial seizures of infancy with quinidine. *Ann Neurol.* 2014;76:457.
- <sup>13</sup>Mikati MA, Jiang Y, Carboni M, et al. Quinidine in the treatment of *KCNT1*-positive epilepsies. *Ann Neurol.* 2015;78:995.
- <sup>14</sup>Milligan CJ, Li M, Gazina EV, et al. *KCNT1* gain of function in 2 epilepsy phenotypes is reversed by quinidine. *Ann Neurol.* 2014;75:581.
- <sup>15</sup>Hoffmann GF, Schmitt B, Windfuhr M, et al. Pyridoxal 5'-phosphate may be curative in early-onset epileptic encephalopathy. *J Inherit Metab Dis.* 2007;30:96.
- <sup>16</sup>Saneto RP, Lee IC, Koenig MK, et al. *POLG* DNA testing as an emerging standard of care before instituting valproic acid therapy for pediatric seizure disorders. *Seizure.* 2010;19(3):140.
- <sup>17</sup>Vigevano F. Benign familial infantile seizures. *Brain Dev.* 2005;27:172.
- <sup>18</sup>Li HF, Chen WJ, Ni W, et al. *PRRT2* mutation correlated with phenotype of paroxysmal kinesigenic dyskinesia and drug response. *Neurology.* 2013;80(16):1534.
- <sup>19</sup>Sheffer IE. Genetic testing in epilepsy: what should you be doing? *Epilepsy Currents.* 2011;11(4):107.
- <sup>20</sup>Wagnon JL, Meisler MH. Recurrent and non-recurrent mutations of *SCN8A* in epileptic encephalopathy. *Front. Neurology.* 2015;6:104.
- <sup>21</sup>Boerma RS, Braun KP, van de Brock M, et al. Remarkable phenytoin sensitivity in 4 children with *SCN8A*-related epilepsy: A molecular neuropharmacological approach. *Neurotherapeutics.* 2016;13:192.
- <sup>22</sup>Barker BS, Ottolini M, Wagnon JL, et al. The *SCN8A* encephalopathy mutation p.Ile1327Val displays elevated sensitivity to the anticonvulsant phenytoin. *Epilepsia.* 2016; published online ahead of print.
- <sup>23</sup>Henske EP, Józwiak S, Kingswood JC, et al. Tuberous sclerosis complex. *Nat Rev Dis Primers.* 2016; published online ahead of print.
- <sup>24</sup>Krueger DA, Wilfong AA, Holland-Bouley K, et al. Everolimus treatment of refractory epilepsy in tuberous sclerosis complex. *Ann Neurol.* 2013;74(5):679.